




Razi University



Original paper

## Optimization of dye removal from aqueous Remazol Brilliant Blue R (RBBR) by *Trametes* sp. Pellets

Parviz Mohamadi<sup>1,2,\*</sup>, Liza Ferina<sup>1</sup>, Mohamad Suffian Mohamad Annuar<sup>3</sup>, Shaliza Ibrahim<sup>2</sup>

<sup>1</sup>Department of Environmental Health Engineering-Kermanshah, Health Research Center (KHRC), Kermanshah University of Medical Science, Kermanshah, Iran.

<sup>2</sup>Department of Civil Engineering, Faculty of Engineering, University of Malaya, 50603 Kuala Lumpur, Malaysia.

<sup>3</sup>Institute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia.

### ARTICLE INFO

#### Article history:

Received 11 August 2013

Received in revised form 29 October 2013

Accepted 14 Feb 2014

Available online 21 March 2014

#### Keywords:

*Trametes* sp.

Remazol Brilliant Blue R

Response surface

Decolorization

### ABSTRACT

Decolorization of Remazol Brilliant Blue R (RBBR) by *Trametes* sp. Pellets was studied in a batch reactor. Dye removal process was performed in shaken flasks which contained 100 ml of RBBR aqueous solution and fungal pellets. The process was followed for 48 h and the dye removal was analyzed at a visible spectrum of 590 nm. Response surface methodology (RSM) employing Box Behnken design at three factors i.e. initial concentration of RBBR, mass of pellet and pH was used to optimize the decolorization process) with three replicates. Response surface regression showed that the decolorization efficiency was affected by initial RBBR concentration. Mass of pellet and pH in this model were not found to be insignificant for both main and square effects. The dye decolorization varied within the range of 16.81% to 77.91%. The lowest decolorization was achieved in maximum initial concentration dye and pH. While the highest decolorization was observed when low initial dye concentration and pH were used. From the optimization, maximum dye removal efficiency of 67.9% ± 5.43 was achieved at 50 ppm RBBR solution, 4 gram of pellets at pH 5.6.

©2014 Razi University-All rights reserved.

### 1. Introduction

Approximately 10,000 different dyes and pigments are manufactured worldwide with the total market of more than 7 x 10<sup>5</sup> tonnes per year. There are several structural varieties of dyes, such as acidic, reactive, basic, disperse, azo, anthraquinone-based, and metal complex dyes (Doble and Kumar 2005). They are used as coloring agents in food, cosmetics, paper, plastic, and textile industries (Cetin and Donmez 2006; Sathiya et al. 2007). Two percent of dyes that are produced by dye industries are discharged directly in the effluent. In addition, 10% is lost during coloration process in textile manufacture (Doble and Kruthiventy 2007). Reactive dyes are widely used in textile industries to colour the cellulosic fibres (Tavares et al. 2009). More than 80,000 tonnes of reactive dyes are produced and consumed every year. This condition described the total amount of pollution generated (Hessel et al. 2007).

Studies have shown that many of the dyes are carcinogenic, mutagenic, and highly harmful to the environment. Untreated dye effluent is highly colored and hence may significantly affect photosynthesis activity due to reduced light penetration (Banat et al. 1996). Several technologies are available for decolorization of textile dye effluents, such as adsorption, irradiation, ion exchange, oxidation, coagulation and precipitation, aerobic process, and anaerobic process. But the problem has not been solved because of high cost, low efficiency, sludge handling problems, less microbial resistant to the pollutant, etc (Anjaneyulu et al. 2005). Recently, many studies of biological decolorization utilizing fungal strains have been reported (Deveci et al. 2004). White rot fungi have been shown to degrade a wide variety of recalcitrant organic pollutant (Young and Yu 2007).

Corresponding author E-mail: [parviz8855@yahoo.com](mailto:parviz8855@yahoo.com)

The several fungal strains were found potential to decolorize commercial reactive dyes e.g. *Bjerkandera adusta*, *Trametes versicolor* and *Phanerochaete chrysosporium* (Heinfling et al. 1997; Swamy and Ramsay 1999).

In this study, *Trametes* sp. pellets were used for decolorizing reactive dye Remazol Brilliant Blue R (RBBR). RBBR is usually used in the production of polymeric dyes. This dye is an anthraquinone derivative which represents a class of toxic and organopollutant materials (Wesenberg et al. 2003). The aim of the studies was to investigate the effect of initial dye concentration, pH, and mass of fungal pellet to decolorization, and optimizing all the variables using response surface methodology to reach the maximum decolorization.

### 2. Materials and methods

#### 2.1. Dye

Reactive Remazol Brilliant Blue R (RBBR) was purchased from Sigma Aldrich, Inc. This dye also have known as Cavalite Brilliant Blue R, C.I. Reactive Blue 19, C.I. Reactive Blue 19 disodium salt, Reactive Blue 19, or Remalan Brilliant Blue R with the empirical formula C<sub>22</sub>H<sub>16</sub>N<sub>2</sub>Na<sub>2</sub>O<sub>11</sub>S<sub>3</sub> and the molecular weight is 626.54. The RBBR was classified as Vinyl Sulfone Reactive dye. Chemical formula of RBBR is presented in Fig. 1.

#### 2.2. Culture

*Trametes* sp. culture was obtained from the Institute of Biological Sciences University of Malaya. The white rot fungus was cultivated on agar medium containing (per liter) 20 g glucose, 0.5 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 1 g K<sub>2</sub>HPO<sub>4</sub>, 0.46 g KH<sub>2</sub>PO<sub>4</sub>, 2 g yeast extract, 2 g malt extract, 2 g

peptone, 0.1 g NH<sub>4</sub>Cl and 18 g agar, and incubated at 28°C for 7 days.




Fig. 1. Remazol Brilliant Blue R.

2.3. Fungal pellet

Fungal pellets were grown in 100 ml liquid medium in 250 ml flask. Culture medium for growing the pellet contained (g/L); 20 glucose, 0.5 MgSO<sub>4</sub>·7H<sub>2</sub>O, 1 K<sub>2</sub>HPO<sub>4</sub>, 0.46 KH<sub>2</sub>PO<sub>4</sub>, 2 yeast extract, 2 malt extract, and 2 peptone. This medium was aseptically inoculated with 5 ml mycelium suspension from actively growing culture on agar plate. Subsequently, this culture was incubated on an orbital shaker (160 rpm) at 28°C for 5 days.

2.4. Response surface design

In this study, response surface employing Box-Behnken design at three factors (initial dye concentration, pH, and mass of pellet) with three replicates was employed. These factors were selected because they were hypothesized to affect decolorization reaction. The levels of the factors are presented in Table 1 and total experimental runs are given in Table 2.

Table 1. Experimental factor and their levels for a three-level Box-Behnken design.

Coded factor	Factors	Coded level		
		-1	0	+1
x <sub>1</sub>	Initial dye concentration	50	65	80
x <sub>2</sub>	mass of pellet	4	5	6
x <sub>3</sub>	pH	4	5	6

The experimental design and analysis of variance (ANOVA) were performed using Minitab® Release 14.12.0 statistical software (Minitab Inc.). A polynomial regression model was used to approximate the response (eq.1).

$$\eta = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_{11} X_1^2 + \beta_{22} X_2^2 + \beta_{33} X_3^2 + \beta_{12} X_1 X_2 + \beta_{13} X_1 X_3 + \beta_{23} X_2 X_3 \tag{1}$$

Where η is the response (decolorization efficiency); X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> are coded levels of the independent factors.

Table 2. Experimental data for RBBR decolorization and percentage of dye removal.

Run	Factors			% decolorization		Run	Factors			% decolorization	
	x <sub>1</sub> (ppm)	x <sub>2</sub>	x <sub>3</sub> (g)	Actual	Predicted		x <sub>1</sub> (ppm)	x <sub>2</sub>	x <sub>3</sub> (g)	Actual	Predicted
1	65	6	6	27.96	29.19	24	65	6	6	25.26	29.19
2	80	5	6	20.40	21.75	25	50	4	5	68.73	63.42
3	50	5	4	59.20	61.78	26	65	4	4	23.47	30.86
4	50	4	5	72.11	63.42	27	65	5	5	30.22	29.22
5	50	5	6	47.45	61.78	28	65	5	5	30.32	29.22
6	65	5	5	29.61	29.22	29	65	6	6	28.66	29.19
7	80	5	6	18.76	21.75	30	80	4	5	20.95	23.39
8	65	6	4	28.59	29.19	31	65	4	6	34.84	30.86
9	80	6	5	21.87	21.72	32	65	5	5	32.26	29.22
10	65	4	4	22.89	30.86	33	50	5	6	53.83	61.78
11	65	6	4	33.30	29.19	34	50	6	5	61.43	61.75
12	65	5	5	30.90	29.22	35	80	5	6	16.81	21.75
13	80	5	4	26.51	21.75	36	50	5	4	77.91	61.78
14	50	5	6	55.94	61.78	37	50	4	5	62.44	63.42
15	80	5	4	27.47	21.75	38	65	4	6	35.77	30.86
16	65	6	4	24.36	29.19	39	65	5	5	30.80	29.22
17	50	5	4	55.98	61.78	40	65	5	5	31.18	29.22
18	50	6	5	71.66	61.75	41	65	5	5	30.79	29.22
19	65	5	5	30.55	29.22	42	65	4	6	32.65	30.86
20	80	6	5	20.22	21.72	43	80	4	5	22.03	23.39
21	65	4	4	28.98	30.86	44	80	5	4	27.29	21.75
22	80	6	5	22.76	21.72	45	80	4	5	20.82	23.39
23	50	6	5	59.57	61.75						

The regression coefficients are: β<sub>0</sub> the constant term; β<sub>1</sub>, β<sub>2</sub>, β<sub>3</sub> the coefficients for linear effects; β<sub>11</sub>, β<sub>22</sub>, β<sub>33</sub> the coefficients for quadratic effects, and β<sub>12</sub>, β<sub>13</sub>, β<sub>23</sub> the coefficient for interaction effects.

2.5. Decolorization studies

The stock solution of RBBR was prepared at 250 ppm and required concentrations were obtained by dilutions. Dye decolorization was performed in Erlenmeyer flasks using the 5-day old pellets at various levels of pH, mass of pellets and initial dye concentration as formulated by the experimental design. The initial pH of aqueous RBBR solution valued 5.6 was adjusted to the desired value using 0.01 M HCl and 0.01 M NaOH.

The flasks were shaken at 160 rpm for 2 days. Samples were withdrawn from the flask, and centrifuged at 7000 rpm for 15 minutes. The absorbance of supernatant was analyzed at a visible spectrum of 590 nm using UV-Vis spectrophotometer (Jasco, Japan). A standard calibration curve was prepared for the dye and the equation is:

$$C_{dye} = 101.83Abs \tag{2}$$

Where C dye is concentration of dye (ppm), and Abs is absorbance. The coefficient of correlation (R<sup>2</sup>) value for above equation is 0.99143. The amount of decolorization was determined using eq. (3).

$$Decolorization(\%) = \frac{(C_i - C_f)}{C_i} \times 100\% \tag{3}$$

Where  $C_i$  is the initial concentration of dye (ppm) and  $C_f$  is the final concentration (ppm).

### 3. Results and discussion

After The results of 45 run for three variables (initial dye concentration, pH, and mass of pellets) for RBBR decolorization are shown in Table 2. The dye decolorization varied within the range of 16.81% to 77.91%. The lowest decolorization was achieved in maximum initial concentration dye and pH. While the highest decolorization was observed when low initial dye concentration and pH were used. Response surface methodology (RSM) was employed to identify the factors that influenced decolorization process, to evaluate interaction that may exist between the factors, and to optimize the factors level to get the maximum decolorization.

A p value is the indicator of the significance of the test, a value below 0.05 indicates that test parameter is significant at 5% level of significance. Analysis of variance (ANOVA) for percentage decolorization which is presented by Table 3 showed that the polynomial regression model is not a fit regression for the data, where the p value of lack of fit is 0.000. It indicates the significant value for lack of fit. Main and square effects are found to be significant ( $p < 0.05$ ), and interaction effects are not significant at 5% level.

Getting a fit regression model, a best subset regression could be employed to find the selection variables for model building as illustrated by Table 4. A good model should have high  $R^2$  and adjusted  $R^2$ , small S, and Mallows' Cp close to the number of variables plus the constant contained in the model. From the Table 4, the Mallows Cp value is closed to row 4 (initial concentration of dye and pH), with  $R^2$  (76.5) and adjusted  $R^2$  (75.4).

To verify the model, the response surface analysis is operated to the selected variables using the linear and square regression model. A new analysis of variance from this model was presented in Table 5. ANOVA shows that fitted linear and square model is highly significant with F-test = 95.08 ( $p = 0.000$ ), and not significant for the lack of fit ( $p = 0.147$ ).

**Table 3.** Analysis of variance for decolorization by polynomial regression model.

Source	DF	Seq SS	Adj SS	Adj MS	F	P
Regression	9	11619.1	11619.1	1291.01	46.89	0.000
Linear	3	9690.1	9690.1	3230.04	117.31	0.000
Square	3	1821.5	1821.5	607.15	22.05	0.000
Interaction	3	107.5	107.5	35.85	1.30	0.289
Residual	35	963.7	963.7	27.53		
Error	3	421.6	421.6	140.53	8.30	0.000
Lack of fit	32	542.1	542.1	16.94		
Pure error						
Total	44	12582.8				

**Table 4.** Best subsets regression: percentage of decolorization versus concentration of dye, mass of pellets.

No.	$R^2$	$R^2$ (adj)	Mallows Cp	S	Initial conc.	Pellets mass	pH
1	76.4	75.9	1.1	8.3086	X		
1	0.5	0.0	136.5	17.066		X	
2	76.9	75.8	2.2	8.3230	X	X	
2	76.5	75.4	2.8	8.3832	X		X
3	77	75.3	4	8.3996	X	X	X

The regression coefficients, t and p values for all the linear, quadratic effects of the variables are given in Table 6. A reduced polynomial was generated relating the response to the significant linear and square regression model i.e:

$$\eta = 29.22 - 20.015X_1 - 0.835X_2 + 12.5467X_1^2 + 0.8117X_2^2 \tag{4}$$

**Table 5.** Analysis of variance for decolorization using the linear and square regression model.

Source	DF	Seq SS	Adj SS	Adj MS	F	P
Regression	4	11385.3	11385.31	2846.33	95.08	0.000
Linear	2	9631.1	9631.14	4815.57	160.85	0.000
Square	2	1754.2	1754.18	877.09	29.30	0.000
Residual Error	40	1197.5	1197.50	29.94		
Lack of fit	4	201.2	201.26	50.29	1.82	0.147
Pure error	36	996.3	996.34	27.68		
Total	44	12582.8				

**Table 6.** Estimated regression coefficients for percentage decolorization of RBBR.

Terms	Coefficients	SE coefficients	t-value	p-value
Constant	29.22	1.518	19.25	0.000
dye concentration	-20.015	1.117	-17.92	0.000
pH	-0.835	1.117	-0.748	0.459
dye concentration*dye concentration	12.5467	1.639	7.655	0.000
pH*pH	0.8117	1.639	0.495	0.623

The coefficient of correlation  $R^2$  for the above equation was 90.48%. Therefore, this equation can be used for predicting response at any combination of three variables within experimental range. The closer the values of R to 1, better the correlation between the experimental and predicted values (Montgomery, 2005). The predicted response value using eq. (4) are shown in Table 2.

For the main factors, only initial dye concentration is found to be significant ( $p < 0.05$ ), where the decolorization percentage decreasing with the initial dye concentration (Fig. 2).

The pH ( $p = 0.459$ ) is found to be insignificant on dye decolorization within the experimental range tested. Meanwhile, for the square effect only initial dye concentration is found to be significant to the decolorization percentage ( $p < 0.05$ ). The square effect of pH ( $p = 0.623$ ) is insignificant on the decolorization of dye.

Response Optimizer function of Minitab was used to find the optimum values of key variables. The solution of eq. (4) is aimed to result in minimum 65% of the dye decolorization. Based on this, a global solution of 50 ppm initial concentration of dye, pH 4 are obtained from the initial point of 50 ppm initial dye concentration, 4 gram pellets mass at pH 4.

The maximum percentage of decolorization predicted by the software was 63.43%, with composite desirability value of 0.96. For practical purposes, the value of the pH is kept at 5.6, respectively in view of the scaling up process.

The slight difference in the value of pH and mass of the pellet was proven to be insignificant to the degree of decolorization as shown by the simulation results where percentage of decolorization was predicted at 61.57%, with composite desirability value of 0.90. Verification experiments were subsequently performed to confirm the

optimization results. The decolorization percentage from these experiments was  $67.9\% \pm 5.4$ .

#### 4. Conclusion

The aqueous RBBR decolorization studies were successfully optimized by response surface methodology. Response surface method explained the effect of the factors and the possible interactions.

The linear and square regression model showed a good fit to the experimental data ( $R^2 = 90.48\%$ ). From the experiment, initial dye concentration was found to be a significant main factor to

decolorization process. After optimization by using statistical analysis, a verification experiment was done and the maximum decolorization efficiency of  $67.9 \pm 5.4\%$  was achieved when optimum value of 50 ppm initial concentration of dye, 4 gram of pellet and pH 5.6 were used.

#### Acknowledgement

The authors would like to acknowledge financial support from the Ministry of Science, Technology and Innovation (03-01-03SF0325), and thank University of Malaya for additional financial support provided through IPPP grant (PS 110-2008C).




Fig. 2. Main effects plot (data means) for percentage of decolorization.

#### Reference

- Anjaneyulu Y., Chary N.S., Raj D.S.S., Decolourization of industrial effluent, available methods and emerging technologies: a review, Review in Environmental Science and Biotechnology 4 (2005) 245-273.
- Banat I.M., Nigam P., Singh D., Marchant R., Microbial decolorization of textile dye containing effluents: a review, Bioresource Technology 58 (1996) 217-227.
- Cetin D., Donmez G., Decolorization of reactive dye by mixed culture isolated from textile effluent under anaerobic conditions, Enzyme and Microbial Technology 38 (2006) 926-930.
- Deveci T., Unyayar A., Mazmanci M.A., Production of Remazol Brilliant Blue R decolourising oxygenase from the culture filtrate of *Funaliatrogii* ATCC 20080, Journal of Molecular Catalyst B: Enzymatic 30 (2004) 25-32.
- Doble M., Kruthiventy A.N., Green chemistry and process, Academic Press, (2007).
- Doble M., Kumar A., Biotreatment of industrial effluents, Butterworth-Heinemann, (2005).
- Heinfling A., Bergbauer M., Szewzyk U., Biodegradation of azo and phthalocyanine dyes by *Trametes* sp. pellets, and *Bjerkandera adusta*, Applied Microbiology and Biotechnology 48 (1997) 261-266.
- Hessel C., Allegre C., Maisseu M., Charbit F., Moulin P., A review Guidelines and legislation for dye house effluents, Journal of Environmental Management 83 (2007) 171-180.
- Montgomery D.C., Design and analysis of experiments, John Wiley and Sons Inc, (2005).
- Sathiya M.P., Periyar S.S., Sasikalaveni A., Murugesan K., Kalaichevan P.T., Decolorization of textile dyes and their effluents using white rot fungi, African Journal of Biotechnology 6 (2007) 424-429.
- Swamy J., Ramsay J.A., The evaluation of white rot fungi in decolorization in the decolorization of textile dyes, Enzyme and Microbial Technology 24 (1999) 130-137.
- Tavares A.P.M., Cristovao R.O., Loureiro J.M., Boaventura R.A.R., Macedo E.A., Application of statistical experimental methodology to optimize reactive dye decolourization by commercial laccase, Journal of Hazardous Materials 162 (2009) 1255-1260.
- Wesenberg D., Kyriakides I., Agathos S.N., White-rot fungi and their enzymes for the treatment of industrial dye effluents, Biotechnology Advance 22 (2003) 61-187.
- Young L., Yu J., Ligninase catalysed decolorization of synthetic dyes, Water Research 31 (2007) 1187-1193.